



SUGGESTED GUIDELINES FOR MANAGEMENT OF CONCUSSION IN SPORTS

National Federation of State High School Associations (NFHS) Sports Medicine Advisory Committee (SMAC)

Introduction

A concussion is type of traumatic brain injury that interferes with normal function of the brain. It occurs when the brain is rocked back and forth or twisted inside the skull as a result of a blow to the head or body. What may appear to be only a mild jolt or blow to the head or body can result in a concussion.

The understanding of sports-related concussion has evolved dramatically in recent years. We now know that young athletes are particularly vulnerable to the effects of a concussion. Once considered little more than a “ding” on the head, it is now understood that a concussion has the potential to result in short or long-term changes in brain function, or in some cases, death.

What is a concussion?

You’ve probably heard the terms “ding” and “bell-ringer.” These terms were once used to refer to minor head injuries and thought to be a normal part of sports. There is no such thing as a minor brain injury. Any suspected concussion must be taken seriously. A concussion is caused by a bump, blow, or jolt to the head or body. Basically, any force that is transmitted to the head causes the brain to literally bounce around or twist within the skull, potentially resulting in a concussion.

It used to be believed that a player had to lose consciousness or be **“knocked-out” to have a concussion.** This is not true, as the vast majority of concussions do not involve a loss of consciousness. In fact, less than 10% of players actually lose consciousness with a concussion.

What exactly happens to the brain during a concussion is not entirely understood. It appears to be a very complex injury affecting both the structure and function of the brain. The sudden movement of the brain causes stretching and tearing of brain cells, damaging the cells and creating chemical changes in the brain. Once this injury occurs, the brain is vulnerable to further injury and very sensitive to any increased stress until it fully recovers.

Common sports injuries such as torn ligaments and broken bones are structural injuries that can be seen on MRIs or x-rays, or detected during an examination. A concussion, however, is primarily an injury that interferes with how the brain works. While there is damage to brain cells, the damage is at a microscopic level and cannot be seen on MRI or CT scans. Therefore, the brain looks normal on these tests, even though it has been seriously injured.

Recognition and Management

If an athlete exhibits any signs, symptoms, or behaviors that make you suspicious that he or she may have had a concussion, that athlete must be removed from all physical activity, including sports and recreation. Continuing to participate in physical activity after a concussion can lead to worsening concussion symptoms, increased risk for further injury, and even death.

SYMPTOMS REPORTED BY ATHLETE
Headache
Nausea
Balance problems or dizziness
Double or fuzzy vision
Sensitivity to light or noise
Feeling sluggish
Feeling foggy or groggy
Concentration or memory problems
Confusion

Parents and coaches are not expected to be able to “diagnose” a concussion. That is the role of an appropriate health-care professional. However, you must be aware of the signs, symptoms and behaviors of a possible concussion, and if you suspect that an athlete may have a concussion, then he or she must be immediately removed from all physical activity.

SIGNS OBSERVED BY PARENTS, FRIENDS, TEACHERS OR COACHES
Appears dazed or stunned
Is confused about what to do
Forgets plays
Is unsure of game, score, or opponent
Moves clumsily
Answers questions slowly
Loses consciousness
Shows behavior or personality changes
Can't recall events prior to hit
Can't recall events after hit

When in doubt, sit them out!